

## Factors influencing lower esophageal sphincter relaxation after deglutition

Lita Tibbling, Per Gezelius, Thomas Franzén

Lita Tibbling, Thomas Franzén, Department of Surgery, University of Linköping, SE-581 85 Linköping, Sweden  
Per Gezelius, Department of Surgery, University of Linköping, SE-581 85; SynMed, SE-117 43 Stockholm, Sweden

Author contributions: Tibbling L, Gezelius P and Franzén T contributed equally to this work; Tibbling L and Franzén T participated in the development, implementation and management of this project and were involved in drafting the manuscript; Gezelius P participated in the analysis of the high-resolution manometry readings and in drafting this part of the manuscript.

Correspondence to: Dr. Thomas Franzén, Department of Surgery, Linköping University Hospital, SE 581 85 Linköping, Sweden. [thomas.franzen@lio.se](mailto:thomas.franzen@lio.se)

Telephone: +46-10-1030000 Fax: +46-10-1043216

Received: December 1, 2010 Revised: March 1, 2011

Accepted: March 8, 2011

Published online: June 21, 2011

**CONCLUSION:** LES relaxation seemed to be caused by the peristaltic wave pushing the bolus from behind against the LES gate.

© 2011 Baishideng. All rights reserved.

**Key words:** Deglutition; Lower esophageal sphincter; Peristalsis; Relaxation; Upper esophageal sphincter

**Peer reviewer:** Kevin Michael Reavis, MD, Assistant Clinic Professor, Department of Surgery, Division of Gastrointestinal Surgery, University of California, Irvine Medical Center, 333 City Boulevard West, Suite 850, Orange, CA, United States

Tibbling L, Gezelius P, Franzén T. Factors influencing lower esophageal sphincter relaxation after deglutition. *World J Gastroenterol* 2011; 17(23): 2844-2847 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v17/i23/2844.htm> DOI: <http://dx.doi.org/10.3748/wjg.v17.i23.2844>

### Abstract

**AIM:** To study the relationship between upper esophageal sphincter (UES) relaxation, peristaltic pressure and lower esophageal sphincter (LES) relaxation following deglutition in non-dysphagic subjects.

**METHODS:** Ten non-dysphagic adult subjects had a high-resolution manometry probe passed transnasally and positioned to cover the UES, the esophageal body and the LES. Ten water swallows in each subject were analyzed for time lag between UES relaxation and LES relaxation, LES pressure at time of UES relaxation, duration of LES relaxation, the distance between the transition level (TL) and the LES, time in seconds that the peristaltic wave was before (negative value) or after the TL when the LES became relaxed, and the maximal peristaltic pressure in the body of the esophagus.

**RESULTS:** Relaxation of the LES occurred on average 3.5 s after the bolus had passed the UES and in most cases when the peristaltic wave front had reached the TL. The LES remained relaxed until the peristaltic wave faded away above the LES.

### INTRODUCTION

Swallowing dysfunction with esophageal food retention is a common problem in an adult population with hiatus hernia. Most articles reporting lower esophageal sphincter (LES) studies depict LES relaxation caused by neurogenic mechanisms<sup>[1]</sup> in gastroesophageal reflux (GER). The ability of the LES to allow esophagogastric transit after deglutition has received little attention. Efforts have been made to prove that LES relaxation is triggered by a neurogenic reflex following deglutition<sup>[2]</sup>, that the LES and the upper esophageal sphincter (UES) relaxations are simultaneous events and that the cervical portion of the vagus nerve mediates inhibitory and excitatory changes in LES pressure<sup>[3]</sup>. In a recent study, a biodynamic approach was proposed for LES opening when GER takes place in hiatal hernia and the gastric wall tension pulls open the LES<sup>[4]</sup>. As long ago as 1978<sup>[5]</sup>, LES was suggested to be a biodynamic gate which is forced to open when a bolus is propelled by esophageal peristalsis. We decided to further

challenge the idea of LES as an esophagogastric pressure zone dependent on complex neuro-humoral factors.

The recent development of high-resolution solid-state manometry (HRM) systems with closely spaced circumferential pressure sensors has made it possible to display simultaneous recordings along the entire esophagus in color-coded pressure plots (Figure 1) and has dramatically improved the diagnostic assessment of esophageal function and disease. With conventional esophageal manometry, simultaneous events in the UES and LES have been difficult to display especially since deglutition often will displace the pressure sensors due to shortening of the esophagus. The HRM technique enables, therefore, a unique possibility to study UES and LES pressure during the entire deglutition period and independent of any sphincter dislocation. In order to find out if there is an interplay between pulling forces and LES opening, the aim was therefore to study the relationship between UES relaxation, peristaltic pressure and LES relaxation following deglutition using the HRM technique.

## MATERIALS AND METHODS

The HRM system (ManoScan 360 A-100, ManoView analysis software ver. 2.0.1 from Sierra Scientific Instruments Inc., Los Angeles, CA) uses a solid state catheter (Ø 4.2 mm) with closely spaced circumferential pressure sensors with 1 cm intervals over 36 cm. The HRM catheter was passed transnasally and positioned to be recording simultaneously from the hypopharynx, through the body of the esophagus, to the stomach. The catheter was calibrated outside the patient before and immediately after the investigation using the thermal compensation option in the software.

We performed a prospective HRM study in 10 adult patients (median age 45 years, range 38-63 years; 7 women, 3 men) who were admitted to an esophageal laboratory for suspected dyspepsia. It can be claimed that our patients with dyspepsia are not representative for studies of normal LES and UES function. Dyspepsia is a diagnosis without organic lesion and with diffuse symptoms predominantly in the gastric region. The patients did not have any symptoms of dysphagia and reflux, they were free of medication, and the presence of hiatus hernia was excluded at gastroscopy and HRM. It is therefore regarded that confounding factors have been excluded in the study material.

The investigation comprised ten swallows of a 10 mL water bolus in a supine position. The characteristics analyzed were: (1) the time lag between UES relaxation and LES relaxation; (2) the LES pressure at time of UES relaxation; (3) the duration of LES relaxation; (4) the length of the esophagus between UES and LES; (5) the distance between the transition level (TL) and the LES; (6) time in seconds that the peristaltic front wave was before (negative value) the TL or after the TL when the LES pressure dropped to nadir; and (7) the maximal peristaltic pressure in the esophagus. The eight best readable swallow recordings in each subject (a total of 78 swallows) were calculated upon, and the mean values of each individual item are given.

TL is defined as an esophageal zone with striated-to-smooth muscle fiber transition<sup>[6]</sup> showing as a short loss of peristaltic pressure (Figure 1).


## RESULTS

Individual values and the mean of all values are presented in Table 1. The LES relaxed on average 3.5 s after the UES had opened. The mean LES pressure was 26 mmHg at time of UES opening. The mean duration of LES remaining relaxed was 6.7 s. The mean length of the esophagus between UES and LES was 26 cm. The mean distance between the TL and the LES was 18 cm, corresponding to on average 72% of the total length. The mean time that the peristaltic wave was before (negative value) the TL or after the TL when the LES opened was -0.5 s. The mean maximum pressure of the peristaltic wave was 138 mmHg. The time lag between UES and LES relaxation was never over 6.4 s. The individual LES pressure, as well as the maximum peristaltic pressure, varied remarkably from swallow to swallow (Table 1).

## DISCUSSION

This study shows quite clearly that the LES becomes relaxed several seconds (on average 3.5 s) after the water bolus has passed the UES. This corresponds with findings by Nguyen *et al.*<sup>[7]</sup> when impedance and manometry techniques were used. They found a mean latency between bolus entry into the esophagus and LES relaxation of 3.6 s. This is also in agreement with the esophageal transit time (3.8 s) as assessed with the biomagnetic technique<sup>[8]</sup>. The LES pressure did not change in our study at time of UES opening. Therefore, it seems unlikely that LES relaxation is triggered by the start of deglutition.

It seems as if opening of the LES coincides with the peristaltic wave front reaching close to the TL, which is an esophageal level with striated-to-smooth muscle fiber transition. Reasonably, this depends on the premise that the distal end of the water bolus has reached the LES, that the water bolus fills up the esophageal lumen between TL and LES, and that the peristaltic wave pushes the water bolus from behind against the LES gate. Only the combination of manometry and impedance measurement or the combination of HRM and radiography can find out whether LES relaxation and opening take place at the moment when the bolus arrives at the LES. In an impedance study of LES opening and bolus transit, it was found that LES relaxation seemed to be modulated by the bolus transit and occurred predominantly upon arrival of the bolus in the distal esophagus<sup>[7]</sup>. This discovery clashes with the findings by Pandolfino *et al.*<sup>[9]</sup> who used a combined impedance/manometry technique. They showed that LES relaxation did not necessary coincide with bolus passage or LES opening. The hydrostatic pressure in the esophageal body necessary to open the LES in an upright position, as shown in a combined manometric and radiographic study from 1978<sup>[5]</sup>, was approximately the same as the LES pressure before opening. If no deglutition activ-



**Figure 1** An high-resolution solid-state manometry recorded swallowing event. Transition level = TL, LES relaxation = LES opening to LES closure. Red  $\geq$  60 mmHg. Blue  $\leq$  0 mmHg. LES fluctuations represent respiration, upwards expiration, downwards inspiration. UES: Upper esophageal sphincter; LES: Lower esophageal sphincter.

Event	F1	F2	M3	F4	F5	F6	M7	M8	F9	F10	All 10 subjects; mean: ranges
(A) s	2.6	5.2	5.0	3.6	3.8	3.2	3.6	1.7	3.0	3.5	3.5: 1.7-5.2
(B) mmHg	21	19	20	32	13	24	23	52	29	29	26: 13-52
(C) s	5.1	4.2	4.2	6.7	5.5	7.7	10.3	8.5	7.7	7.5	6.7: 4.2-10.3
(D) cm	25	26	28	25	29	23	26	26	22	26	26: 22-29
(E) cm	16	17	20	18	20	17	19	20	16	18	18: 16-20
(F) s	-1.4	0.7	0.9	0.3	0.5	-0.2	-1.4	-2.0	-1.6	-0.3	-0.5: -2.0-0.9
(G) mmHg	109	121	123	71	102	232	187	169	152	116	138: 71-232

Different esophageal events, mean values of ten recordings in ten subjects, and mean values of all ten subjects. A: Time lag between relaxing of UES and LES; B: LES pressure when UES opened; C: Duration of LES remaining relaxed; D: Esophagus length; E: Distance between TL and LES; F: Time in seconds that the peristaltic wave was before (minus value) or after the TL when the LES relaxed; G: maximum pressure of the peristaltic wave. F: Female; M: Male. UES: Upper esophageal sphincter; LES: Lower esophageal sphincter; TL: Transition level defined as the zone when the striated muscle layer transitions into the smooth muscle layer.

ity took place and a contrast medium was instilled into the esophagus, the LES opened when the hydrostatic pressure exceeded the resting LES pressure and closed again when the hydrostatic pressure in an upright position fell short of the LES pressure. Deglutition caused the LES to relax and open when the peristaltic wave reached the upper level of the infused contrast medium. These different findings indicate that the LES is a barrier which is forced to open by the peristaltic pressure. In the clinic, this would explain why, for instance, patients with achalasia cardia or with lack of esophageal peristalsis do not have any LES relaxation after deglutition.

If we look upon the LES as a gate that will be pulled open, it is of interest to compare the pressure of the LES

and the maximum pressure of the peristaltic wave in the distal esophagus. In this study, the peristaltic pressure was five times stronger than the LES pressure. HRM and conventional manometry are claimed to be the same in their measurement of LES resting pressure<sup>[10]</sup>. It has been shown that the longitudinal esophageal muscle is contracted during the peristaltic activity<sup>[11]</sup> which will decrease esophageal wall compliance. Certainly, the peristaltic force displays an interaction between the longitudinal and circular esophageal muscles<sup>[12]</sup>, and a compliance decrease of the esophageal wall will facilitate bolus transit. The reason for the deglutition-induced pressure overload in the esophageal body is therefore of importance, in order to overcome the stiffness of the esophageal wall. In the study by Babaei *et al*<sup>[12]</sup>, it was

even proposed that the longitudinal esophageal smooth muscle has an important role in the relaxation of LES. In our study, the LES remained relaxed as long as the peristaltic wave was present in the esophagus. This indicates that there is a close interplay between LES function and peristaltic activity. It is reasonable to believe that the LES will be pulled open when the pulling direction is either from the esophagus or from the stomach<sup>[12]</sup>.

In conclusion, LES opening seems to be caused by the peristaltic wave pushing the bolus from behind against the LES gate. Therapeutic attention in patients with dysphagia of non-stricture origin should therefore be focused on esophageal motility function rather than on drugs affecting LES pressure.

## COMMENTS

### Background

Swallowing problems with food retention in the gullet are present in at least 8% of an adult population. For accurate treatment, it is of importance to know whether transit from the esophagus to the stomach can be treated with drugs aimed at opening the lower esophageal sphincter (LES) or whether transit is due to dynamic properties of gullet muscles. So far, it has been claimed that LES is triggered to open by a neurogenic reflex from the upper esophageal sphincter (UES).

### Research frontiers

The newly developed high-resolution manometry (HRM) system is a technical innovation and breakthrough for the understanding of dynamic esophageal events, meaning the interplay between esophageal motility and esophageal sphincter relaxations.

### Innovations and breakthroughs

The esophageal HRM probe was used in ten non-dysphagic patients with dyspepsia in order to study the time relationship between UES and LES relaxation and to study where the peristaltic wave front was located when the LES relaxed after deglutition of a 10 mL water bolus in the supine position. These simultaneous activities have previously not been possible to study with conventional esophageal manometry.

### Applications

The LES was shown to relax 3.5 s later than the UES after deglutition which is the average time it takes a water bolus front to reach the LES. The LES remained relaxed on average 6.8 s; that is until the propulsive force had faded away. The peristaltic front wave reached a level close to the transition level (TL), either 2 s before or 1 s after the TL.

### Terminology

Esophagus-gullet. Dyspepsia-a diagnosis without any specified organic lesion and with diffuse symptoms from the gastric region. Dysphagia-swallowing difficulties. HRM-a pressure catheter with a 36 cm long segment of sensors spaced 1 cm apart giving simultaneous pressure information of sphincter and muscular activity from a total of 432 locations from UES to LES. The pressure can be displayed as a color plot offering visual information of the esophagus and its sphincters very similar to an anatomic picture. LES-lower esophageal sphincter. Manometry-pressure measurement. Smooth muscle-a muscle under autonomic, non-volitional control. Striated muscle-a muscle under volitional control. TL, transition level-the

level in the esophagus between the upper striated muscle level and the lower smooth muscle level which is located about 7 cm distal of the UES.

### Peer review

More emphasis on why HRM is a great technique and the knowledge it provides compared to conventional esophageal manometry has now been given in the Methods section. The reason why we believe that dyspepsia patients can be regarded as normal people with regard to esophageal transit studies is addressed in the Materials section. The previous conclusion, regarding older theories of a neurogenic reflex causing LES relaxation, has been omitted and basic data for this are given in the discussion.

## REFERENCES

- 1 Tuch A, Cohen S. Lower esophageal sphincter relaxation: studies on the neurogenic inhibitory mechanism. *J Clin Invest* 1973; **52**: 14-20
- 2 Bardan E, Saeian K, Xie P, Ren J, Kern M, Dua K, Shaker R. Effect of pharyngeal stimulation on the motor function of the esophagus and its sphincters. *Laryngoscope* 1999; **109**: 437-441
- 3 Matarazzo SA, Snape WJ Jr, Ryan JP, Cohen S. Relationship of cervical and abdominal vagal activity to lower esophageal sphincter function. *Gastroenterology* 1976; **71**: 999-1003
- 4 Fein M, Ritter MP, DeMeester TR, Oberg S, Peters JH, Hagen JA, Bremner CG. Role of the lower esophageal sphincter and hiatal hernia in the pathogenesis of gastroesophageal reflux disease. *J Gastrointest Surg* 1999; **3**: 405-410
- 5 Ask P, Sökjer H, Tibbling L. Mechanisms affecting lower oesophageal sphincter opening and oesophageal retention. A combined X-ray and manometry study. *Scand J Gastroenterol* 1978; **13**: 857-861
- 6 Ghosh SK, Janiak P, Schwizer W, Hebbard GS, Brasseur JG. Physiology of the esophageal pressure transition zone: separate contraction waves above and below. *Am J Physiol Gastrointest Liver Physiol* 2006; **290**: G568-G576
- 7 Nguyen HN, Domingues GR, Winograd R, Lammert F, Silny J, Matern S. Relationship between bolus transit and LES-relaxation studied with concurrent impedance and manometry. *Hepatogastroenterology* 2006; **53**: 218-223
- 8 Daghananli NA, Braga FJ, Oliveira RB, Baffa O. Oesophageal transit time evaluated by a biomagnetic method. *Physiol Meas* 1998; **19**: 413-420
- 9 Pandolfino JE, Shi G, Zhang Q, Ghosh S, Brasseur JG, Kahrilas PJ. Measuring EGJ opening patterns using high resolution intraluminal impedance. *Neurogastroenterol Motil* 2005; **17**: 200-206
- 10 Ayazi S, Hagen JA, Zehetner J, Ross O, Wu C, Oezcelik A, Abate E, Sohn HJ, Banki F, Lipham JC, DeMeester SR, Demeester TR. The value of high-resolution manometry in the assessment of the resting characteristics of the lower esophageal sphincter. *J Gastrointest Surg* 2009; **13**: 2113-2120
- 11 Tibbling L, Ask P, Pope CE 2nd. Electromyography of human oesophageal smooth muscle. *Scand J Gastroenterol* 1986; **21**: 559-567
- 12 Babaei A, Bhargava V, Korsapati H, Zheng WH, Mittal RK. A unique longitudinal muscle contraction pattern associated with transient lower esophageal sphincter relaxation. *Gastroenterology* 2008; **134**: 1322-1331

S- Editor Sun H L- Editor Logan S E- Editor Ma WH